


AD-8167 484 ON L SUB 1-CONTRACTION FOR SYSTEMS OF CONSERVATION LAWS 1/1  
(U) WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER  
J G PATINO FEB 86 MRC-TSR-2917 DAG29-80-C-0041

UNCLASSIFIED

F/G 12/1

NL





MICROCOPY

CHART

MRC Technical Summary Report #2917

ON L<sub>1</sub>-CONTRACTION FOR SYSTEMS OF  
CONSERVATION LAWS

Jorge G. S. Patiño

AD-A167 484

Mathematics Research Center  
University of Wisconsin—Madison  
610 Walnut Street  
Madison, Wisconsin 53705

February 1986

(Received February 24, 1986)

DTIC  
ELECTE  
MAY 23 1986  
S D

Approved for public release  
Distribution unlimited

Sponsored by

U. S. Army Research Office  
P. O. Box 12211  
Research Triangle Park  
North Carolina 27709

86 5 20 136

DTIC FILE COPY

UNIVERSITY OF WISCONSIN-MADISON  
MATHEMATICS RESEARCH CENTER

ON  $L_1$ -CONTRACTION FOR SYSTEMS OF CONSERVATION LAWS<sup>1</sup>

Jorge G. S. Patiño

Technical Summary Report #2917  
February 1986

ABSTRACT

We prove that for  $2 \times 2$ , strictly hyperbolic, genuinely nonlinear systems of conservation laws, there is no metric  $D$  such that

$$\int_{-\infty}^{\infty} D(u(x,t), c) dx$$

is a decreasing function of time for every weak solution  $u$ ,  $u_0(\pm\infty) = c$ .

AMS (MOS) Subject Classifications: 35L65, 76N10

Key Words: Conservation laws, Decay of solutions,  $L_1$ -dependence

Work Unit Number 1 (Applied Analysis)

---

<sup>1</sup>Work done while the author was a visiting member of the Mathematics Research Center, University of Wisconsin-Madison.

Present address: Departamento de Matematica, PUC-RJ, Rua Marquês de São Vicente 225, 22453 Rio de Janeiro.

---

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041.

# SIGNIFICANCE AND EXPLANATION

The Cauchy problem for a  $2 \times 2$  system of conservation laws in one dimension is

$$u_t + (f(u))_x = 0, \quad x \in \mathbb{R}, \quad t > 0$$

$$u(x, 0) = u_0(x)$$

where  $u = (u_1, u_2)$ ,  $f = (f_1(u), f_2(u))$ .

Such systems of equations usually come from the application of the laws of conservation for physical quantities like mass, momentum and energy, and arise in problems of gas dynamics, elasticity, oil reservoir simulation and other areas of engineering.

The questions of decay and continuous dependence with respect to the initial data are central issues in the study of the problem above. The result proved here rules out the use of certain functionals to study the decay of solutions and is relevant to the issue of  $L_1$  continuity with respect to the data.

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	



The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

# ON $L_1$ -CONTRACTION FOR SYSTEMS OF CONSERVATION LAWS<sup>1</sup>

Jorge G. S. Patiño

For  $2 \times 2$ , strictly hyperbolic, genuinely nonlinear (cf. [1]) systems of conservation laws it was proved in [2] that there is no metric  $D$ , compatible with the state space, such that

$$I_D(u, v; t) \equiv \int_{-\infty}^{\infty} D(u(x, t), v(x, t)) dx \quad (1)$$

is a decreasing function of time for any two weak solutions  $u, v$  whose initial conditions agree off a compact set.

In [2] a metric  $D$  is compatible with the state space  $\Sigma$  if

C1.  $D : \Sigma \times \Sigma \rightarrow \mathbb{R}$

is a symmetric function.

C2.  $D(u, v) + D(v, w) \geq D(u, w) \quad \forall u, v, w \in \Sigma$

C3.  $C_0^{-1}|u - v| \leq D(u, v) \leq C_0|u - v| \quad \forall u, v \in \Sigma$

with a uniform constant  $C_0$ .

Here we give an easier proof of this result, and give conditions that a metric must satisfy if condition C3 is to be relaxed.

We wish to point out that relaxing condition C3 is important since it rules out the use e.g. of entropies or quadratic functions to obtain certain integral decay estimates. It is also interesting to note that the solutions used in the

---

<sup>1</sup>Work done while the author was a visiting member of the Mathematics Research Center, University of Wisconsin-Madison.

Present address: Departamento de Matematica, PUC-RJ, Rua Marquês de São Vicente 225, 22453 Rio de Janeiro.

construction of the counter examples below are the elementary "spikes" used frequently in decay arguments (cf. [3]).

Thus let

$$u_t + (f(u))_x = 0 \quad (2)$$

be any  $2 \times 2$  system, strictly hyperbolic and genuinely nonlinear on a region  $N \subset \mathbb{R}^2$ . Let  $\lambda_1(u)$  and  $\lambda_2(u)$  be the eigenvalues of  $df(u)$  with corresponding eigenvectors  $r_1(u)$  and  $r_2(u)$ . Let  $R_1(u, u^*)$  and  $R_2(u, u^*)$  be respectively the integral curves of  $r_1(u)$  and  $r_2(u)$  passing through  $u^*$ .  $R_1(u, u^*)$  and  $R_2(u, u^*)$  are called rarefaction curves. Let  $S_1(u, u^*)$  and  $S_2(u, u^*)$  be the curves of states that can be joined by respectively a 1-shock and 2-shock to the right of  $u^*$ . These are called shock curves. Given a state  $u^*$  on  $N$ , shock and rarefaction curves exist locally [1].

We then have the following theorem

Theorem 1. Let  $u$  and  $v$  be weak solutions of (2) whose initial conditions agree off a compact set, then there exists no metric  $D$ , which is symmetric, such that  $I_D(u, v; t)$  is a strictly decreasing function of time.

Proof. Take any states  $u_L, u_R, \bar{u}$  and  $\bar{\bar{u}}$  related in the following way

(Figure 1)

- i)  $u_R$  and  $u_L$  are joined by a 1-shock with speed  $s_1$ , with  $u_L$  on the left.
- ii)  $u_R$  and  $\bar{u}$  are joined by a 1-rarefaction.
- iii)  $\bar{u}$  and  $u_L$  are joined by a 2-rarefaction.
- iv)  $\bar{\bar{u}}$  and  $u_L$  are joined by a 2-rarefaction.
- v)  $\bar{\bar{u}}$  and  $u_L$  are joined by a 1-rarefaction.

(We assume here that  $\lambda_2(u)$  increases from  $\bar{u}$  to  $u_L$ . The case where  $\lambda_2(u)$  decreases from  $\bar{u}$  to  $u_L$  is discussed below.)




Figure 1

The system (2) with initial condition

$$u(x,0) = \begin{cases} u_R & \text{if } 0 < x < (s_1 - \lambda_1(u_R))T \\ u_L & \text{otherwise} \end{cases}$$

has, for  $t \leq T$ , the solution  $u$  shown in Figure 2.




Figure 2. A line denotes a shock and a fan denotes a rarefaction.

Then

$$I_D(u, u_L, 0) = D(u_R, u_L)(s_1 - \lambda_1(u_R))T$$

and

$$I_D(u, u_L, T) = T \int_{\lambda_1(u_R)}^{\lambda_1(\bar{u})} D(u(\lambda), u_L) d\lambda + D(u_L, \bar{u})(\lambda_2(\bar{u}) - \lambda_1(\bar{u}))T \\ + T \int_{\lambda_2(\bar{u})}^{\lambda_2(u_L)} D(u(\lambda), u_L) d\lambda ,$$



where  $\mu(\lambda)$  and  $v(\lambda)$  denote parametrizations of  $R_1(u, \bar{u})$  with respect to  $\lambda_1$  and of  $R_2(u, u_L)$  with respect to  $\lambda_2$ , respectively.

Now, with  $u_R, u_L$  and  $\bar{u}$  denoting the same states as in Figure 1, consider the following initial condition:

$$v(x, 0) = \begin{cases} u_L & \text{if } 0 < x < (\lambda_2(u_L) - s_1)T \\ u_R & \text{otherwise.} \end{cases}$$

The solution  $v$  of this problem, for  $t < T$ , is given by the waves in Figure 3.




Figure 3

Then

$$I_D(v, u_R; 0) = D(u_L, u_R)(\lambda_2(u_L) - s_1)T$$

and

$$I_D(v, u_R; 0) = T \int_{\lambda_1(u_R)}^{\lambda_1(\bar{u})} D(\mu(\lambda), u_R) d\lambda + T \int_{\lambda_2(\bar{u})}^{\lambda_2(u_L)} D(v(\lambda), u_R) d\lambda \\ + D(\bar{u}, u_R)(\lambda_2(\bar{u}) - \lambda_1(\bar{u}))T.$$

To prove the theorem by contradiction, assume now that

$$I_D(v, u_R; T) + I_D(u, u_L; T) < I_D(v, u_R; 0) + I_D(u, u_L; 0). \quad (3)$$

Thus

$$\begin{aligned}
& \lambda_1(\bar{u}) \\
& \int_{\lambda_1(u_R)}^{\lambda_1(\bar{u})} \{D(\mu(\lambda), u_L) + D(\mu(\lambda), u_R)\} d\lambda \\
& + \int_{\lambda_2(\bar{u})}^{\lambda_2(u_L)} \{D(v(\lambda), u_L) + D(v(\lambda), u_R)\} d\lambda \\
& + D(u_L, \bar{u})(\lambda_2(\bar{u}) - \lambda_1(\bar{u})) + D(\bar{u}, u_R)(\lambda_2(\bar{u}) - \lambda_1(\bar{u})) \\
& \leq D(u_R, u_L)(s_1 - \lambda_1(u_R)) + D(u_L, u_R)(\lambda_2(u_L) - s_1) .
\end{aligned}$$

Now, adding and subtracting

$$D(u_L, u_R)(\lambda_1(\bar{u}) - \lambda_1(u_R)) + D(u_L, u_R)(\lambda_2(u_L) - \lambda_2(\bar{u}))$$

we get

$$\begin{aligned}
& \lambda_1(\bar{u}) \\
& \int_{\lambda_1(u_R)}^{\lambda_1(\bar{u})} \{D(\mu(\lambda), u_L) + D(\mu(\lambda), u_R) - D(u_L, u_R)\} d\lambda \\
& + \int_{\lambda_2(\bar{u})}^{\lambda_2(u_L)} \{D(v(\lambda), u_L) + D(v(\lambda), u_R) - D(u_R, u_L)\} d\lambda \quad (4) \\
& + (D(u_L, \bar{u}) + D(\bar{u}, u_R) - D(u_R, u_L))(\lambda_2(\bar{u}) - \lambda_1(\bar{u})) \leq 0 .
\end{aligned}$$

By the triangle inequality the two integrands and the third line above are positive. Since  $\lambda_1(u_R) < \lambda_1(\bar{u})$  and  $\lambda_2(\bar{u}) < \lambda_2(u_L)$ , equality holds in (4) if and only if equality holds in each of the triangle inequalities, in particular only if

$$D(u_L, \bar{u}) + D(\bar{u}, u_R) = D(u_R, u_L) . \quad (5)$$

A similar construction as above using the initial conditions

$$u_0(x) = \begin{cases} u_L & \text{if } 0 < x < \lambda_1(u_L) - s \\ \bar{u} & \text{otherwise} \end{cases}$$

and

$$u_0^1(x) = \begin{cases} \bar{u} & \text{if } 0 < x < \lambda_2(v) - \lambda_1(v) \\ u_L & \text{otherwise} \end{cases}$$

will, by the same argument, yield

$$D(u_R, \bar{u}) + D(u_R, u_L) = D(\bar{u}, u_L) . \quad (6)$$

Now, (5) and (6) give

$$D(\bar{u}, u_L) - D(\bar{u}, u_R) = D(u_L, \bar{u}) + D(u_R, \bar{u}) . \quad (7)$$

This equation gives, using the compatibility condition C3. Lemma 4 in [2] after which the proof proceeds identically.

If one does not assume C3, (5) and (6) put restrictions on the metric  $D$ . For instance the Euclidean distance or a linear combination of changes in Riemann invariants will not work.

In the case where  $\lambda_2(u)$  decreases from  $\bar{u}$  to  $u_L$  a similar construction using the shock curves  $S_2(u, \bar{u})$  and  $S_2(u, \hat{u})$  (see Figure 4) and the same initial value problems as above yield conditions similar to (5) and (6) from where the proof would proceed identically.




Figure 4

Acknowledgments. I would like to thank Bradley Plohr for reading the first draft and for a good suggestion. I also thank Blake Temple for many discussions on relevant issues. This work was made possible by the financial support of the Conselho Nacional de Pesquisa-CNPq, Brazil and the Financiadora de Estudos e Projetos-FINEP, Brazil.

#### REFERENCES

1. Lax, P. D., Hyperbolic systems of conservation laws II., Comm. Pure Appl. Math. 19 (1957), 537-566.
2. Temple, J. B., No  $L_1$ -contractive metrics for systems of conservation laws, Trans. Amer. Math. Soc. 288-2 (1985), 471-480.
3. Temple, J. B., Decay with a rate for noncompactly supported solutions of conservation laws, Trans. Amer. Math. Soc., to appear.

JGSP:scr

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 2917	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ON $L_1$ -CONTRACTION FOR SYSTEMS OF CONSERVATION LAWS		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Jorge G. S. Patiño		8. CONTRACT OR GRANT NUMBER(s) DAAG29-80-C-0041
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of Wisconsin 610 Walnut Street Madison, Wisconsin 53705		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 1 - Applied Analysis
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P.O. Box 12211 Research Triangle Park, North Carolina 27709		12. REPORT DATE February 1986
		13. NUMBER OF PAGES 7
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)  UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Conservation laws Decay of Solutions $L_1$ -dependence		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) We prove that for $2 \times 2$ , strictly hyperbolic, genuinely nonlinear systems of conservation laws, there is no metric $D$ such that $\int_{-\infty}^{\infty} D(u(x,t), c) dx$ is a decreasing function of time for every weak solution $u$ , $u_0(\pm\infty) = c$ .		

END

DTIC

6-86